

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 7 : C07C 235/26, 237/18, 235/30, 237/20, C07D 211/76, 265/24, 401/06, A61K 31/4402, 31/45, 31/4545, 31/535, A61P 9/10, 3/06, 29/00, 37/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/48989 (43) International Publication Date: 24 August 2000 (24.08.00) <i>27</i></p>
<p>(21) International Application Number: PCT/EP00/01191 (22) International Filing Date: 14 February 2000 (14.02.00) (30) Priority Data: 9903546.1 16 February 1999 (16.02.99) GB 9927880.6 25 November 1999 (25.11.99) GB (71) Applicant (for all designated States except AT US): NOVAR- TIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN [AT/AT]; Verwaltungsgesellschaft m.b.H., Brunner Strasse 59, A-1230 Vienna (AT). (72) Inventors; and (75) Inventors/Applicants (for US only): BAENTELI, Rolf [CH/CH]; Sennheimerstrasse 51, CH-4054 Basel (CH). BAUER, Wilfried [CH/CH]; Hohli Gasse 7; CH-4432 Lampenberg (CH). COTTENS, Sylvain [CH/CH]; Trauben- weg 34, CH-4108 Witterswil (CH). EHRHARDT, Claus [DE/DE]; Grossmannstrasse 17, D-79541 Lörrach (DE). HOMMEL, Ulrich [DE/DE]; Vögisheimer Weg</p>		<p>9, D-79379 Müllheim (DE). KALLEN, Jörg [CH/CH]; Kaltbrunnenstrasse 41, CH-4054 Basel (CH). MEIN- GASSNER, Josef, Gottfried [AT/AT]; Max Margules Weg 10, A-2380 Perchtoldsdorf (AT). NUNINGER, François [FR/FR]; 11, rue de l'Entente, F-68270 Wittenheim (FR). WEITZ-SCHMIDT, Gabriele [DE/DE]; Steinbrecherstrasse 4, D-79189 Bad-Krozingen (DE). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, ER, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GI, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: MEVINOLIN DERIVATIVES (57) Abstract Mevinolin derivatives wherein the lactone ring is modified have interesting pharmaceutical properties, particularly in preventing or treating disorders or diseases mediated by LFA-1/ICAM-1 interactions.</p>		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Mevinolin Derivatives

The present invention relates to mevinolin derivatives, a process for their production, their use as a pharmaceutical and pharmaceutical preparations containing them.

More particularly the present invention provides a compound of formula I



wherein

each of a—b and α—β independently, is either a single bond or a double bond;


R₁ is



wherein R_a is H; C₁₋₆alkyl optionally substituted by OH or C₁₋₄alkoxy; C₂₋₆alkenyl; or aryl-C₁₋₄alkyl;

R₂ is OH; -O-CO-R₅ wherein R₅ is C₁₋₈alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkyl, aryl or aryl-C₁₋₄alkyl; or -O-R₆ wherein R₆ is the residue of an α-amino-acid attached to O through its carbonyl residue or -CH(R₇)-COR₈ wherein R₇ is H, C₁₋₄alkyl, heteroC₁₋₄alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkyl, aryl or aryl-C₁₋₄alkyl and R₈ is OH, C₁₋₄alkoxy or NR₉R₁₀ wherein each of R₉ and R₁₀ independently is H, C₁₋₄alkyl or R₉ and R₁₀ form together with the nitrogen to which they are bound, a heteroaryl group;

R₃ is a substituted lactam, piperidyl, linear amino alcohol or cyclic carbamate, or a residue of formula (i)



wherein

R_{13} is OH; C_{1-6} alkoxy; $-O-CO-C_{1-6}$ alkyl; or $-O-CO-NHC_{1-6}$ alkyl;

R_{14} is OH; C_{1-4} alkoxy; C_{1-4} alkyl; C_{1-4} alkoxy-carbonyl- C_{1-4} alkoxy; hydroxy- C_{1-5} alkoxy; C_{1-4} alkoxy- C_{1-5} alkoxy; C_{1-4} alkoxy-carbonyl- C_{1-4} alkyl; or $NR_{9a}R_{10a}-C_{1-5}$ alkoxy wherein each of R_{9a} and R_{10a} independently has one of the significances given for R_9 and R_{10} ;

R_{15} is H or C_{1-4} alkyl; and

R_{16} is $CONR_{17}R_{18}$ wherein one of R_{17} and R_{18} is H and the other is C_{1-6} alkyl, hydroxy- C_{1-6} alkyl, C_{3-7} cycloalkyl- C_{1-4} alkyl or aryl- C_{1-4} alkyl; or C_{1-6} alkoxy-carbonyl; each of $\alpha-b$ and $\alpha-\beta$ being a single bond when each of R_{13} or R_{14} is OH; and

R_4 is H or OR_{19} wherein R_{19} is C_{1-6} alkyl, hydroxy- C_{1-6} alkyl, C_{1-4} alkoxy- C_{1-6} alkyl, aryl- C_{1-4} alkyl or C_{1-4} alkoxycarbonyl- C_{1-4} alkyl,

and wherever "aryl" appears as is or in the significances of "aryl- C_{1-4} alkyl" in the above definition, it is "phenyl" or "naphthyl" optionally substituted by halogen, OH, $NR_{11}R_{12}$, COOH, CF_3 , C_{1-4} alkoxy, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, hydroxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl, cyano or $CONR_{11}R_{12}$, each of R_{11} and R_{12} independently being H, C_{1-4} alkyl, phenyl, naphthyl, phenyl- C_{1-4} alkyl or naphthyl- C_{1-4} alkyl or R_{11} and R_{12} together with the nitrogen to which they are bound forming heteroaryl; and wherever "heteroaryl" appears, it is a 5- or 6-membered heterocyclic residue optionally fused to a benzene ring; in free form or in salt form.

Alkyl groups or alkyl moieties may be branched or straight chain. Cycloalkyl groups or moieties are preferably cyclopentyl or cyclohexyl. Heteroalkyl includes e.g. halogenated alkyl such as CF_3 . Polyhydroxy- C_{1-4} alkyl may comprise up to 6 hydroxy groups.

Preferably the phenyl or naphthyl moiety in aryl or aryl- C_{1-4} alkyl, when substituted, bears up to 3 substituents as disclosed above, more preferably selected from C_{1-4} alkoxy, e.g. methoxy or ethoxy, hydroxy- C_{1-4} alkoxy, hydroxy- C_{1-4} alkyl and OH. When the phenyl moiety is disubstituted, the 2 substituents are preferably in positions meta and para. Aryl- C_{1-4} alkyl is preferably benzyl, phenethyl or naphthyl- CH_2- , the phenyl or naphthyl moiety being optionally substituted as indicated above.


Examples of heteroaryl include pyrrolyl, imidazolyl, furyl, thienyl, pyrrolidinyl, piperidyl, piperazinyl, morpholino, pyridyl, indolyl or quinolyl. Heteroaryl as formed by R_9 and R_{10} together with the nitrogen to which they are attached, may comprise a further heteroatom.

e.g. O or N, and is preferably pyrrolidinyl, piperidyl, piperazinyl or morpholino. In heteroaryl-C₁₋₄alkyl, the alkyl moiety preferably is C₁ or C₂alkyl.

The significances given above for "aryl" and "heteroaryl" also applies to the radicals of formulae (a), (b), (c₁) or (c₂) hereinafter.

When R₆ is the residue of an α-amino acid, it may be the residue of a natural or unnatural α-amino acid residue, e.g. Ala, Leu, Ile, Val, Pro, wherein the terminal amino group may be substituted or unsubstituted, e.g. by an amino protecting group.

When R₃ is a substituted lactam residue, it is preferably a 6-membered ring wherein the nitrogen of the lactam may be substituted and/or comprising a further substituent on the ring, e.g. on the carbon atom opposite to the nitrogen. Preferably the lactam residue is disubstituted. A suitable example of a substituted lactam as R₃ includes e.g. a radical of formula (a)




wherein

R₃₀ is C₁₋₆alkyl; C₃₋₇cycloalkyl; aryl; C₃₋₇cycloalkyl-C₁₋₄alkyl; aryl-C₁₋₄alkyl; heteroaryl; or heteroaryl-C₁₋₄alkyl;

R₃₁ is OH; C₁₋₄alkoxy; C₁₋₄alkyl; C₁₋₄alkoxy-carbonyl-C₁₋₄alkoxy; hydroxy-C₁₋₅alkoxy; C₁₋₄alkoxy-C₁₋₅alkoxy; C₁₋₄alkoxy-carbonyl-C₁₋₄alkyl; amino-C₁₋₄alkoxy;

HOOC-C₁₋₄alkoxy; HOOC-C₁₋₄alkyl; or NR_{9a}R_{10a}-C₁₋₅alkoxy wherein each of R_{9a} and R_{10a} independently has one of the significances given for R₉ and R₁₀.

When R₃ is a substituted piperidyl residue, the nitrogen of the piperidyl may be substituted and/or a further substituent may be present on the ring, e.g. on the carbon atom opposite to the nitrogen. Preferably the piperidyl residue is disubstituted. A suitable example of a substituted piperidyl residue includes e.g. a radical of formula (b)




wherein

R_{40} has one of the significances given for R_{30} ; and

R_{41} has one of the significances given for R_{31} or is $-O-CO-C_{1-8}$ alkyl.

When R_3 is a substituted amino alcohol residue, the amino group thereof may be monosubstituted, e.g. by a substituent such as aryl- C_{1-4} alkyl or aryl- C_{1-4} alkyl-carbonyl, and/or a further substituent may be present on the chain, e.g. on the carbon atom adjacent to the alcohol or amino group. Cyclisation of the substituted amino alcohol residue leads to a corresponding substituted cyclic carbamate. A suitable example of a substituted amino alcohol and of the corresponding substituted cyclic carbamate includes e.g. a radical of formula (c₁) or (c₂)



wherein

either each of X and Y is H or X and Y form together



each of R_{50} , independently is H; C_{1-8} alkyl; C_{3-7} cycloalkyl; aryl; C_{3-7} cycloalkyl- C_{1-4} alkyl; aryl- C_{1-4} alkyl; heteroaryl; heteroaryl- C_{1-4} alkyl; C_{1-4} alkyl-carbonyl; aryl-carbonyl; heteroaryl-carbonyl; aryl- C_{1-4} alkyl-carbonyl or heteroaryl- C_{1-4} alkyl-carbonyl, and

each of R_{51} , independently is H; C_{1-4} alkyl; hydroxy- C_{1-4} alkyl; amino- C_{1-4} alkyl; C_{1-4} alkoxy- C_{1-4} alkyl; C_{1-4} alkoxy-carbonyl- C_{1-4} alkyl wherein C_{1-4} alkoxy is optionally substituted by amino, C_{1-4} alkyl-amino or di- $(C_{1-4}$ alkyl)-amino; $HOOC-C_{1-4}$ alkyl; or $R_{23}R_{24}N-CO-C_{1-4}$ alkyl wherein R_{23} is H, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, polyhydroxy- C_{1-8} alkyl, heteroaryl, heteroaryl- C_{1-4} alkyl, amino- C_{1-4} alkyl, C_{1-4} alkylamino- C_{1-4} alkyl, di- $(C_{1-4}$ alkyl)amino-

C_{1-4} alkyl or aryl- C_{1-4} alkyl and R_{24} is H, C_{1-4} alkyl or hydroxy- C_{1-4} alkyl, at least one of R_{50} and R_{51} being other than H.


Preferred compounds of formula I are those wherein R_3 is substituted lactam, substituted linear amino alcohol, substituted cyclic carbamate, preferably substituted lactam or substituted cyclic carbamate, e.g. as disclosed above, more preferably a radical of formula (a) or (c₁) or (c₂) wherein X and Y are -CO-.

In the compounds of formula I, the following significances are preferred individually or in any sub-combination:

1. R_1 is H or CH_3 , preferably CH_3 ;
2. R_2 is -O-CO- R_5 , preferably wherein R_5 is C_{4-8} alkyl, particularly -CH(CH_3)- CH_2 - CH_3 , -CH(CH_2 - CH_2 - CH_3)₂, -CH(CH_2 - CH_3)₂, -C(CH_3)₂- CH_2 - CH_3 or -CH(CH_2 - CH_3)- CH_2 - CH_2 - CH_3 .
3. a--b is a double bond;
4. α -- β is a double bond;
5. R_4 is H;
6. R_3 is a radical of formula (I);
7. R_{16} is CO-NR₁₇R₁₈; preferably one of R_{17} and R_{18} is H;
8. Each of R_{13} and R_{14} is OH and each of a--b and α -- β is a single bond;
9. Each of R_{13} and R_{14} is other than OH;
10. R_3 is a radical of formula (a);
11. R_{30} in (a) is aryl- C_{1-4} alkyl or heteroaryl- C_{1-4} alkyl, preferably benzyl or naphthyl-methyl wherein the phenyl or naphthyl ring is optionally substituted by OH, C_{1-4} alkoxy, hydroxy- C_{1-4} alkoxy or hydroxy- C_{1-4} alkyl, or morpholino, pyridyl, indolyl or quinolyl;
12. R_{31} in (a) is OH, C_{1-4} alkoxy, hydroxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl- C_{1-4} alkoxy or HOOC- C_{1-4} alkoxy;
13. R_3 is a radical of formula (c₁) or (c₂) wherein X and Y form together -CO-;
14. R_{50} in (c₁) or (c₂) wherein X and Y form together -CO-, is aryl- C_{1-4} alkyl or heteroaryl- C_{1-4} alkyl, preferably benzyl or naphthyl-methyl wherein the phenyl or naphthyl ring is optionally substituted by OH, C_{1-4} alkoxy, hydroxy- C_{1-4} alkoxy or hydroxy- C_{1-4} alkyl;
15. R_{51} in (c₁) or (c₂) wherein X and Y form together -CO-, is hydroxy- C_{1-4} alkyl; amino- C_{1-4} alkyl; C_{1-4} alkoxy- C_{1-4} alkyl; C_{1-4} alkoxy-carbonyl- C_{1-4} alkyl; HOOC- C_{1-4} alkyl; or $R_{23}R_{24}N$ -CO- C_{1-4} alkyl.

Compounds of formula I may exist in free form or in salt form, e.g. as acid addition salts with e.g. organic or inorganic acids, for example, hydrochlorides, or salts when a COOH is present, as salts with bases e.g. alkali salts such as sodium or potassium, or substituted or unsubstituted ammonium salts.


It will be appreciated that the radicals of formulae (i), (a), (b), (c₁) and (c₂) may comprise at least one asymmetric carbon atom, e.g. the carbon atom which bears R₁₅ and R₁₆, R₃₁, R₄₁ or R₅₁, respectively, for example




Where the stereochemistry of any part of a compound of the invention is not specified, it is to be understood that the present invention embraces individual enantiomers and their mixtures. Similar considerations apply in relation to starting materials exhibiting asymmetric carbon atoms as mentioned above. Where compounds of the invention exist in isomeric form as aforementioned, individual isomers may be obtained in conventional manner, e.g. employing optically active starting materials or by separation of initially obtained mixtures, for example using conventional chromatographic techniques.

The present invention also includes a process for the production of a compound of formula I, comprising

- a) for the production of a compound of formula I wherein R₃ is a residue of formula (i) submitting mevinolin or compactin or the corresponding tetrahydro-mevinolin or -compactin to ring opening, e.g. by reaction with a corresponding amine, e.g. arylamine; or
- b) for the production of a compound of formula I wherein R₃ is a radical of formula (c₁) wherein each of X and Y is H, submitting to reductive amination the carbonyl function in R₃ in a compound of formula IV



wherein R_1 , R_2 , R_4 , $a-b$ and $\alpha-\beta$ and R_1 are as defined above, and R_3 is a radical of formula (c_{1A})



wherein R_{51} is as defined above; or

- c) for the production of a compound of formula I wherein R_3 is a residue of formula (c₂) wherein each of X and Y is H, submitting mevinolin or compactin wherein the lactone ring has been converted into a conjugated α,β unsaturated lactone, to a 1,4-addition e.g. with an amine, e.g. veratrylamine, and concomitant ring opening with an alcohol, e.g. methanol; or
- d) for the production of a compound of formula I wherein R_3 is a residue of formula (c₁) or (c₂) wherein each of X and Y is -CO-, submitting to cyclisation a compound of formula I wherein R_3 is a residue of formula (c₁) or (c₂) wherein each of X and Y is H; or
- e) for the production of a compound of formula I wherein R_3 is a substituted lactam, e.g. a residue of formula (a), submitting a compound of formula I wherein R_3 is a residue of formula (i) wherein R_{13} is OH oxidised to a ketone and R_{16} is CONHR₁₈, to a reductive amination and concomitant ring closure; or converting the free OH group in R_3 in a compound of formula I wherein R_3 is a residue of formula (i) wherein R_{18} is CONHR₁₈, into a leaving group, e.g. by mesylation, and then submitting the resulting compound to a basic treatment; or
- f) for the production of a compound of formula I wherein R_3 is a substituted piperidyl, e.g. a residue of formula (b), reducing a compound of formula I wherein R_3 is a substituted lactam, e.g. a residue of formula (a);

and, where required, removing the protecting group where present, and converting the resulting compound of formula I in free form or in salt form.

Where OH groups are present in the starting products which are not to participate in the reaction, they may be protected, in accordance with known methods. OH protecting groups are known in the art, e.g. *tert*.-butyl-dimethyl-silyl.

Process steps (a) to (f) may be effected analogously to methods known in the art or as disclosed in the Examples below. The cyclisation in step (d) may conveniently be carried out in the presence of a cyclisation agent, e.g. carbonyl diimidazole.

Compounds of formula IV may be prepared by opening of the OH protected lactone ring according to known procedures, e.g. by reaction with an amine and then oxidation of the resulting hydroxy group into a ketone. Insofar as the production of the starting materials is not particularly described, the compounds are known or may be prepared analogously to methods known in the art or as disclosed in WO 99/11258, e.g. starting from mevinolin or compactin or tetrahydro-mevinolin or -compactin. The -O-CO-CH(CH₃)-C₂H₅ of mevinolin, compactin or tetrahydro-mevinolin or -compactin may also be reduced to OH and then esterified to another -O-CO-R₅ group.

The following Examples are illustrative of the invention. Following abbreviations are used:

Boc = *tert*.-butoxy-carbonyl

rt = room temperature

OMe = methoxy

THF = tetrahydrofuran

DMF = dimethylformamide

DCC = N,N'-dicyclohexylcarbodiimide

Pro = proline

TBDMS = *tert*-butyldimethylsilyl


DMAP = dimethylaminopyridine

CDI = carbonyldiimidazole

TBME = *tert*-butylmethylether

CHX = cyclohexane

Example 1: (S)-2-Methyl-butyric acid (S)-(3R,7S,8aR)-8-[(S)-2-[(4R,6R)-3-(4-hydroxy-3-methoxy-benzyl)-4-[(2-hydroxy-ethylcarbamoyl)-methyl]-2-oxo-[1,3]oxazin-6-yl]-ethyl]-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester



- a) To a solution of 40.4 g mevinolin ((S)-2-Methyl-butyric acid (1S,3R,7S,8S,8aR)-8-[2-((1R,3R)-4-hydroxy-6-oxo-tetrahydro-pyran-2-yl)-ethyl]-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester) in 100 ml of CH_2Cl_2 are added 24.4 g DMAP, then slowly 15.0 g Ac_2O . The mixture is stirred overnight at rt. The reaction is controlled by TLC, TBME/CHX, 3:2. The reaction mixture is diluted with TBME, washed successively with water, ca. 15% citric acid, brine, then dried over sodium sulfate. The organic phase is concentrated and the product crystallized by addition of diisopropylether. The precipitate is filtered, washed with diisopropylether and dried, yielding the α,β unsaturated lactone derivative (S)-2-methyl-butyric acid (S)-(3R,7S,8aR)-7-methyl-3-methyl-8-[(S)-2-((R)-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester.

MS (FAB-MS), 387 (MH⁺)

- b) To 15.5 g of compound a) in 250 ml MeOH is added 14.4 g of the 4-(*tert*-butyl-dimethyl-silanyloxy)-3-methoxy-benzylamine and the mixture is stirred overnight. TLC control in TBME. The reaction mixture is completely evaporated and the crude product separated by flash-chromatography on silica gel (CHX-> TBME-> MeOH). The desired methylester (S)-2-methyl-butyric acid (S)-(3R,7S,8aR)-8-((R)-(3R,5R)-5-[4-(*tert*-butyl-dimethyl-silanyloxy)-3-methoxy-benzylamino]-3-hydroxy-6-methylcarbamoyl-hexyl)-7-methyl-3-methyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester is obtained.

MS, (ESI), 686.5 (MH⁺)


- c) A solution of 12.5 g of the compound as obtained in b), in 25 ml DMF is treated with 4.1 g CDI and stirred for ca. 5 h. at rt. TLC control in TBME/ CHX, 3:2. The reaction mixture is diluted with TBME, extracted with water and then brine, the organic phase dried over sodium sulfate and evaporated. The crude product is purified by RP-18 chromatography, MeOH/H₂O -> MeOH. After rechromatography on silica gel, TBME/CHX -> TBME, the cyclic carbamate (S)-2-Methyl-butyrac acid (S)-(3R,7S,8aR)-8-((S)-2-((4R,6R)-3-[4-(*tert*-butyl-dimethyl-silanyloxy)-3-methoxy-benzyl]-4-methylcarbamoylmethyl-2-oxo-[1,3]oxazinan-6-yl]-ethyl)-7-methyl-3-methyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester is obtained as a foam.

MS (ESI), 712.5 (MH⁺)

- d) To a solution of 6.0 g of the compound as obtained in c) in 25 ml MeOH is added a total of 5.5 g 2-amino-ethanol and the mixture is heated at reflux, for ca. 40 hrs., until completion (TLC control in moist ethyl acetate). The reaction mixture is diluted with ethyl acetate, extracted with citric acid and brine, then dried over sodium sulfate and evaporated. The crude product is purified as above, first by RP-18 chromatography then on silica gel. The title product is obtained as a white foam.

MS (ESI): 627.4 (MH⁺)

Example 2: (S)-2-Methyl-butyrac acid (S)-(3R,7S,8aR)-8-((S)-2-((4R,6R)-3-(4-hydroxy-3-methoxy-benzyl)-4-methylcarbamoylmethyl-2-oxo-[1,3]oxazinan-6-yl]-ethyl)-7-methyl-3-methyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester




3.6 g of the compound as obtained in Example 1c) is dissolved in 100 ml of a solution of ca. 30% methylamine in MeOH and stirred at rt for ca. 24 hrs. (TLC control in moist ethyl

acetate). The reaction mixture is evaporated and the crude product purified by silica gel chromatography (TMBE/CHX → ethyl acetate). Pure fractions are combined, evaporated and the title compound is obtained as a foam.


MS (ESI): 597.4 (MH⁺) $[\alpha]_D^{20} = +234.7^\circ$ (c=1 in methanol)

By following the same procedure, but using the appropriate starting material, the diastereomer wherein the carbamate ring has the configuration



is also obtained.


By following the procedure as disclosed in Examples 1 and 2, the compounds of formula X




wherein R₅₀ and R₅₁ are as defined in Table 1 below, may be prepared.

Table 1

Ex	R ₅₀	R ₅₁	M.S. (ESI)
3*	3-OMe-4-OH-benzyl	-CH ₂ -CO-NH-CH ₂ -(4'-OH-3'-OMe-phenyl)	717 [M-H]
4	3-OMe-4-(2-hydroxy-ethoxy)-benzyl	-CH ₂ -CO-NH-CH ₂ -[3'-OMe-4'-(β-hydroxy-ethoxy)-phenyl]	851 [M+HCOO-]
5	3,4-di-OMe-benzyl	-CH ₂ -CO-NHCH ₃	611 [MH ⁺]
6	3,4-di-OMe-benzyl	-CH ₂ -COOH	596 [M-H]

7	3-OMe-4-OH-benzyl	-CH ₂ -CO-OCH ₃	596 [M-H]
8	3-OMe-4-OH-benzyl	-CH ₂ -COOH	582 [M-H]
9	3-OMe-4-OH-benzyl	-CH ₂ -CO-N(CH ₃) ₂	609 [M-H]
10	3-OMe-4-OH-benzyl	-CH ₂ -CH ₂ -OH	570 [MH+]
11	3,4-di-OH-benzyl	-CH ₂ -CO-NH-CH ₂ -(3',4'-di-OH-phenyl)	689 [M-H]
12	3,4-di-OMe-benzyl	-CH ₂ -CO-N-CH(CH ₂ OH) ₂	669 [M-H]
13	3-OMe-4-(2-hydroxy-ethoxy)-benzyl	-CH ₂ -CO-NHCH ₃	639 [M-H]
14	3-OMe-4-OH-benzyl	-CH ₂ -CO-NH-CH ₂ -CH(OH)-CH ₂ OH	655 [M-H]
15	3,4-di-OH-benzyl	-CH ₂ -CO-OCH ₃	582 [M-H]
16	3,4-di-OH-benzyl	-CH ₂ -CO-NHCH ₂ CH ₂ OH	611 [M-H]
17	3-OMe-4-OH-benzyl	-CH ₂ -CO-NH-CH(CH ₂ OH) ₂	655 [M-H]
18	3,4-di-OMe-benzyl	-CH ₂ -CO-NH-CH ₂ -CH ₂ OH	640 [M-H]
19		-CH ₂ -CO-NH-CH ₂ -CH ₂ OH	689 [M-H]
20	3-OMe-4-OH-benzyl	-CH ₂ -CO-NH-CH ₂ -(CHOH) ₄ -CH ₂ OH	745 [M-H]
21	3-OMe-4-OH-benzyl	-CH ₂ -CO-N-(CH ₂ CH ₂ OH) ₂	669 [M-H]
22	3-OMe-4-OH-benzyl	-CH ₂ -CO-NH-(CH ₂) ₂ -N(CH ₃) ₂	654 [MH+]
23	4-OMe-3-OH-benzyl	-CH ₂ -CO-NH-CH ₂ -CH ₂ OH	671 [M+HCOO-]
24	3,4-di-OMe-benzyl	-CH ₂ -CO-NH-(3,4-di-OMe-benzyl)	721 [MH+]


* The diastereoisomer of the compound of Ex. 3, wherein the cyclic carbamate residue has the configuration



is also prepared by following the same procedure.

By following the procedure below the compounds of formula X₁ may be prepared. The OH protected lactone ring of mevinolin or compactin may also be submitted to ring opening, e.g.

by reaction with an amine, then treatment of the resulting hydroxyamine with carbonyl-diimodazole leads to the carbamate.




wherein R_{50} , R_{51} are as defined in Table 2 below.

Table 2

Ex	R_{50}	R_{51}	M.S. (ESI)
25*	3,4-di-OMe-benzyl	$-\text{CH}_2\text{-CO-NHCH}_3$	611 (M+H)
26	3,4-di-OMe-benzyl	$-\text{CH}_2\text{-CO-O-(CH}_2)_2\text{-N(CH}_3)_2$	668 (M+H)


* The diastereomer of the compound of Ex. 25, wherein the cyclic carbamate residue has the configuration



is also prepared by following the same procedure.


Example 27: (S)-2-Methyl-butyrlic acid (S)-(3S,4aS,7S,5S,8S,8aS)-8-[(S)-2-[(4R,6R)-3-(4-hydroxy-3-methoxy-benzyl)-4-[(2-hydroxy-ethylcarbamoyl)-methyl]-2-oxo-[1,3]oxazinan-6-yl]-ethyl]-3,7-dimethyl-decahydro-naphthalen-1-yl ester

By following the first step of the procedure to prepare example 28 (to obtain trans tetrahydro mevinolin) and then the procedure (step a) to d) as disclosed in Example 1, the compound of formula



is obtained. MS (ESI): 629 [M-H].

Example 28: (S)-2-Methyl-butyrlic acid (S)-(3S,4aS,7S,8S,8aS)-8-[(3R,5R)-6-(3,4-dimethoxy-benzylcarbamoyl)-3,5-dihydroxy-hexyl]-3,7-dimethyl-decahydro-naphthalen-1-yl ester




- a) To a solution of 40 g (0.098 mol) of mevinolin in 3 l ethyl acetate is added 10 g $\text{P}/\text{Al}_2\text{O}_3$. The mixture is hydrogenated under an H_2 atmosphere under 2.6 bar pressure for 16 h. The mixture is filtered and the solvent evaporated. The residue is purified by silical gel chromatography using ethyl acetate/cyclohexane 8/2 as a solvent. First eluted is the undesired cis isomer, followed by a side product with one double bond. Finally the desired trans isomer is eluted. Several crystallizations yielded the desired trans tetrahydro-mevinolin ((S)-2-Methyl-butyrlic acid (S)-(3S,4aS,7S,5S,8S,8aS)-8-[2-((2R,4R)-4-hydroxy-6-oxo-tetrahydro-pyran-2-yl)-ethyl]-3-methyl-7-methyl-decahydro-naphthalen-1-yl ester).

- b) To a solution of 2 g (5.0 mmol) of the trans tetrahydro mevinolin obtained in a) in 12 ml ethanol is added 3.7 ml (25.0 mmol) 3,4-dimethoxybenzylamine. The reaction mixture is stirred for 20 h at rt, then it is diluted with 300 ml diethyl ether with and washed with 100 ml of water. The organic extract is dried with MgSO_4 and evaporated. The residue is purified by chromatography on silicagel using ethyl acetate as eluent to give the title compound.

MS (ESI): 598 (M+Na), 574 (M-H)


The compound of Example 29 is obtained by silylating mevinolin, following step b) of the procedure of Example 28, reacting with ethyl isocyanate and desilylating according to conventional methods. The compound of example 30 is obtained by reacting mevinolin with ethyl diazoacetate and rhodium acetate and then following step b) of the procedure of Example 28.

Example 29:



MS(ESI): 613(M+H)

Example 30:





MS (ESI): 658 (M+H)


The compound of Example 31 is prepared by following step a) and b) of the procedure of Example 1 and desilylating the compound resulting from step b). The compounds of Examples 32 and 33 are obtained by submitting the appropriate starting materials to step a) and a modified version of step b) (in the absence of methanol):

Example 31:

Examples 32 and 33:

MS (ESI): 571 [MH⁺]Ex. 32 R = H MS (ESI): 693 [MH⁺]Ex. 33 R = CH₃ MS (ESI): 791 [M+HCOO]

Example 34: (S)-2-Methyl-butyric acid (1S,3R,7S,8S,8aR)-8-[2-[(2S,4R)-4-hydroxy-1-(5-hydroxymethyl-6-methoxy-naphthalen-2-ylmethyl)-6-oxo-piperidin-2-yl]-ethyl]-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester



- a) To a stirred solution of 22 g (37 mmol) of silylated mevinolin (obtained by standard silylation of mevinolin in the 4 position) ((S)-2-Methyl-butyric acid (3R,7S,8S,8aR)-8-[2-[(2R,4R)-4-(*tert*-butyl-dimethyl-silanyloxy)-6-oxo-tetrahydro-pyran-2-yl]-ethyl]-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester) in 65 ml THF at rt are added 18 g (56 mmol) of C-[5-(*tert*-butyl-dimethyl-silanyloxymethyl)-6-methoxy-naphthalen-2-yl]-methylamine (prepared from 2-bromo-6-methoxy-naphthalene). After 18 hours the reaction mixture is diluted with 250 ml methyl-*t*-butyl ether and washed successively with 10% aqueous citric acid, saturated aqueous sodium bicarbonate and brine. The organic phase is dried over sodium sulfate and the solvent evaporated. The crude product is purified by silica gel chromatography (hexane/ethyl acetate 4/1 to 3/2) to afford the hydroxyamide 2-methyl-butyric acid 8-(5-(*tert*-butyl-dimethyl-silanyloxy)-6-[[5-(*tert*-butyl-dimethyl-silanyloxymethyl)-6-methoxy-naphthalen-2-ylmethyl]-carbamoyl]-3-

hydroxy-hexyl)-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester as a white foam.

MS (ESI, -Q1MS) 894.6; 884.3; 848.5

- b) To a stirred, cooled (0 °C) solution of 4.3 g (5.0 mmol) of the compound obtained under 34a) above and 1.4 ml (10 mmol) triethylamine in 40 ml THF are added 0.51 ml (6.6 mmol) of methanesulfonyl chloride. After 30 minutes 6.5 ml (13 mmol) of a 2M solution of sodium bis(trimethylsilyl)amide in THF are added. The mixture is stirred for 1 hour at 0 °C. the reaction is quenched with 10% aqueous citric acid and diluted with methyl-t-butyl ether. The phases are separated and the aqueous phase is extracted twice with methyl-t-butyl ether. The organic phases are combined, washed successively with saturated aqueous sodium bicarbonate and brine, dried over sodium sulfate and the solvent is evaporated. The crude product is purified by silica gel chromatography (hexane/ethyl acetate 95/5 to 4/1) to afford the lactam 2-methyl-butyric acid 8-(2-{4-(*tert*-butyl-dimethyl-silanyloxy)-1-[5-(*tert*-butyl-dimethyl-silanyloxymethyl)-6-methoxy-naphthalen-2-yl)methyl]-6-oxo-piperidin-2-yl}-ethyl)-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester as a white foam.

MS (ESI, +Q1MS) 832.6; 598.4


- c) To a stirred solution of 106 mg (0.13 mmol) of the compound obtained under 34b) above in 2 ml THF at room temperature are added 606 µl (0.62 mmol) of a 1N aqueous HCl solution. After 18 hours the reaction is quenched with saturated aqueous sodium bicarbonate and diluted with methyl-t-butyl ether. The phases are separated and the aqueous phase is extracted twice with methyl-t-butyl ether. The combined organic phases are washed with brine, dried over sodium sulfate and the solvent is evaporated. The crude product is purified by silica gel chromatography (hexane:ethyl acetate 1:1 to 1:4) to afford the pure title compound as a white foam.

MS (ESI, -Q1MS) 648.4; 602.5

$[\alpha]_D^{20} = +119.3^\circ$ (c=1 in methanol)

m.p. = 145°C


The compounds of formula X₂




wherein R_{30} and R_{31} have the significances given in Table 3, are prepared analogously to the procedure as disclosed in Example 34.

Table 3

Ex	R_{30}	R_{31}	M.S
35		OH	545 [MH+]
36*		OH	545 [MH+]
37	4-(2'-O-1-ethoxy)-3-OCH ₃ -benzyl	OH	628 [M+HCOO-]
38	3,5-di-(OCH ₃)-benzyl	OH	554 [MH+]
39	3,4-di-(OCH ₂ CH ₃)-phenyl	OH	568 [MH+]
40	β-naphthyl-CH ₂ -	OH	544 [MH+]
41	4-di-ethyl-carbamoyl-benzyl	OH	593 [MH+]
42	3-OCH ₃ -4-OH-benzyl	OH	538 [M-H]
43	4-morpholinocarbonyl-benzyl	OH	607 [MH+]
44		OH	619 [M+HCOO-]
45	4-ethoxy-carbonyl-benzyl	OH	566 [MH+]
46*		OH	519 [MH+]
47**	(3,4-dimethoxy)-benzyl	OH	554 [M+H]
48	4-pyridyl-CH ₂ -	OH	495 (M+H)
49**	3-pyridyl-CH ₂ -	OH	494 (M) (EI)
50	CH ₃	OH	418 (M+H)
51	3-benzoyloxy-benzyl	OH	600 (M+H)
52*	3-benzoyloxy-benzyl	OH	600 (M+H)
53	3-isopropoxy-benzyl	OH	596 (M+HCOO)

54*	3-isopropoxy-benzyl	OH	552 (M+H)
55	(3,4-dimethoxy)-phenethyl	OH	568 (M+H)
56	p-CF ₃ -benzyl	OH	562 (M+H)
57	p-tert.-butoxy-benzyl	OH	594 (M+H)
58		OCH ₃	640 [MNa+]
59**	m-methoxy-benzyl	OH	523 (M) (EI)
60*	benzyl	OCH ₃	508 (M+H)
61	3-OCH ₃ -4-(2'-OH-ethoxy)-benzyl	O-CH ₂ -CO-OC ₂ H ₅	715 [M+HCOO-]
62	3-OCH ₃ -4-(2'-OH-ethoxy)-benzyl	O-CH ₂ -CH ₂ -OH	628 [MH+]
63*	benzyl	OCH ₂ CH ₂ OH	538 (M+H)
64*	benzyl	O-CH ₂ -COOH	552 (M+H)

* In these compounds the lactam moiety has the following configuration:




**Both diastereoisomers are obtained and each can be isolated.


The synthesis of compounds of Examples 61 to 64* additionally comprise a treatment with ethyl diazoacetate and rhodium acetate, followed for the compounds of Examples 62* and 63*, by a reduction.

By following the procedure of Example 34 but using as starting material the corresponding tetrahydro-mevinolin derivative and 3,4-dimethoxy-benzylamine, the following compound is obtained:


Example 65:

MS (ESI): 558 [MH⁺]

Compounds of Ex. 66 and 67 may be obtained from mevinolin as follows: ester cleavage of mevinolin and oxidation of the newly generated hydroxy position to the oxo compound. The neighbouring hydroxy substituent is then introduced via the formation of the silylenolate and treatment with meta-chloroperbenzoic acid. Selective alkylation of the newly formed hydroxy position is achieved by treatment with Meerwein salt. The ester group is introduced via its anhydride. Then the procedure as described for Ex. 34 is followed.

Example 66

MS (ESI) 598 (M+H)

Example 67


MS (ESI) 598 (M+H)

Example 68: (S)-2-Methyl-butyrlic acid (1S,3R,7S,8S,8aR)-8-[2-((R)-1-benzyl-4-methyl-6-oxo-piperidin-2-yl)-ethyl]-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester

Mevinolin is treated with acetic anhydride to give the α,β -unsaturated lactone. This is treated with cuprous bromide dimethylsulfide complex and methyl lithium to affect conjugate addition. The methylated lactone compound is treated with methanol and diazabicycloundecane to give the ring opened methylated hydroxy ester. The hydroxy group of which is then oxidized with sulfur trioxide pyridine complex to the corresponding ketone.

The ketone is reductively aminated (as described for Example 76b) to give the title compound. MS(EI): 491 (M)

By following the procedure of Example 68, but using the appropriate starting materials, the compounds of formula X₃



wherein R₁ and Y-Z are as defined in Table 4 below, may be prepared:

Table 4

Example	Y-Z	R ₁	MS (EI)
68	C—C	-CH ₂ -phenyl	see above
69	C=C	-CH ₂ -CH(CH ₃) ₂	457 (M)
70	C≡C	-CH ₂ -CH(CH ₃) ₂	457 (M)

Example 71: (S)-2-Methyl-butyric acid (1S,3R,7S,8S,8aR)-8-{2-[(2S,4S)-1-(3,4-dimethoxy-benzyl)-4-hydroxy-6-oxo-piperidin-2-yl]-ethyl}-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester


a) To a stirred solution of 600 mg (1.1 mmol) of 2-methyl-butyric acid 8-{2-[1-(3,4-dimethoxy-benzyl)-4-hydroxy-6-oxo-piperidin-2-yl]-ethyl}-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester in 15 ml of THF at rt are added 2 ml of a 65% RedAl® solution in toluene. After 3 hours, the reaction is quenched by the addition of 1 ml of methanol. The organic phase is extracted twice with 15 ml of 2N HCl. The aqueous phases are combined, brought to pH 12 with 1N NaOH and extracted three times with ethyl acetate. The organic phases are combined, dried over sodium sulfate and the solvent evaporated. The residue is purified by silica gel chromatography (t-butyl methyl

ether/methanol 9/1) to afford pure 1-(3,4-Dimethoxy-benzyl)-2-[2-(8-hydroxy-2,6-dimethyl-1,2,6,7,8,8a-hexahydro-naphthalen-1-yl)-ethyl]-piperidin-4-ol as a white foam.
MS (ESI) 456 (M+H)

b) The above compound is treated with diethylacetic anhydride in the presence of catalytic amounts of 4-dimethylaminopyridine in dichloromethane at rt for 16 h to give a diacylated compound. The undesired acyl group on the lactone moiety is cleaved by transesterification with methanol at 55°C for 5 h to give the title compound.

MS (ESI): 554 (M+H). See formula below in Table 5.

By following the procedure of Example 71, but using the appropriate starting materials, the compounds of formula X₄



wherein R₁-R₃ and Y-Z are as defined in Table 5 below, may be prepared

Table 5

Ex	R ₁	R ₂	R ₃	Y-Z	MS (ESI)
71	OH	3,4-dimethoxy-benzyl	(CH ₃ -CH ₂) ₂ CH-CO-	C=C	see above
72	OH	3,4-dimethoxy-benzyl	(CH ₃ -CH ₂) ₂ CH-CO-	C≡C	554 (M+H)
73	(CH ₃ -CH ₂) ₂ CH-CO-O-	3,4-dimethoxy-benzyl	(CH ₃ -CH ₂) ₂ CH-CO-	C=C	674 (M+Na)
74	(CH ₃ -CH ₂) ₂ CH-CO-O-	3,4-dimethoxy-benzyl	(CH ₃ -CH ₂) ₂ CH-CO-	C≡C	652 (M+H)
75	(CH ₃ -CH ₂) ₂ CH-CO-O-	3,4-dimethoxy-benzyl	H	C≡C	554 (M+H)

Compound of Ex. 73 is obtained from compound of Ex. 71 starting from compound of Ex. 47. Compound of Ex. 74 is obtained from compound of Ex. 72 starting from the diastereoisomer of Ex. 50**.

Example 76: (S)-2-Methyl-butyric acid (1S,3R,7S,8S,8aR)-8-[(3R,5R)-3-(3,4-dimethoxybenzylamino)-5-hydroxy-6-methylcarbamoyl-hexyl]-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester


a) To a solution of 900 mg (1.65 mmol) of 2-methyl-butyric acid 8-[5-(tert.-butyl-dimethyl-silanyloxy)-6-methylcarbamoyl-3-oxo-hexyl]-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester in 5 ml of THF are added 700 mg (11.7 mmol) of acetic acid and 1.0 g (3.2 mmol) of tetrabutylammonium fluoride trihydrate. The reaction mixture is stirred for 3 hours at rt. It is then diluted with 30 ml of ethyl acetate and washed successively with a saturated aqueous sodium bicarbonate solution and water. The organic phase is dried over sodium sulfate and the solvent evaporated. The residue is crystallized from diethyl ether to afford the desired product 2-methyl-butyric acid 8-(5-hydroxy-6-methylcarbamoyl-3-oxo-hexyl)-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester as white crystals.

MS (FAB) 440 (M+Li)

b) To a solution of 300 mg (0.69 mmol) of the compound of formula 76a) in 2 ml of dichloroethane are added 200 mg (1.2 mmol) of veratrylamine, 244 mg (1.15 mmol) of sodium triacetoxyborohydride and 60 mg (1.0 mmol) of acetic acid. The reaction mixture is stirred overnight at room temperature. It is then diluted with 20 ml of ethyl acetate and washed successively with a saturated aqueous sodium bicarbonate solution, water and brine. The organic phase is dried over sodium sulfate and the solvent evaporated. The residue is purified by silica gel chromatography (tert.-butyl methyl ether/methanol/NH₄OH_{aq} 90/9/1) to afford the title compound as a white foam.

MS (ESI) 585 (M+H)

By following the procedure of Example 76, but using the appropriate starting materials, the compounds of formula X₅



wherein R_1 and R_2 are as defined in Table 6 below, may be prepared.

Table 6

Ex	R_1	R_2	MS (ESI)
76	CH_3	-(3,4-dimethoxy)-benzyl	see above
77	CH_3	benzyl	525 (M+H)
78	CH_3	diphenyl-methyl	601 (M+H)
79	CH_3	$CH(CH_3)$ -phenyl	539 (M+H)

The compounds of formula I, in free form or in pharmaceutically acceptable salt form exhibit valuable pharmacological properties, e.g. inhibiting activity of LFA-1/ICAM-1 or ICAM-3 interactions or inhibiting inflammation, e.g. as indicated in in vitro and in vivo tests and are therefore indicated for therapy.

A. In vitro:

i) Cell Free Assay

The assay measures the binding of soluble human ICAM-1 to immobilized human LFA-1. LFA-1 is purified from JY cells, a human lymphoblastoid B cell-line, by immunoaffinity chromatography as described by Dustin *et al.* (J. Immunol. 148, 2654-2663, 1992). ICAM-1 mouse Cx fusion protein (ICAM-1) is produced using the baculovirus system as described by Weltz-Schmidt *et al.* (Anal. Biochem. 238, 184-190, 1996).

Purified LFA-1 is diluted 1:20 in phosphate buffered saline (PBS) containing 2 mM MgCl_2 , pH 7.4 and coated onto microtitre plates (Nunc) at 37°C for 3h. Plates are blocked with 1% heat-treated BSA in PBS for 2 hours at 37°C followed by a washing step using PBS, 2mM MgCl_2 , 1% fetal calf serum, pH 7.4 (assay buffer). Compounds dissolved at 10 mM in DMSO are diluted in assay buffer and added to the plates. Biotinylated recombinant ICAM-1 in assay buffer (6 $\mu\text{g/ml}$) is added and allowed to bind at 37°C for one hour. After incubation, wells are washed with assay buffer. Streptavidin-peroxidase diluted 1:5000 in assay buffer is added and incubated for 45 min at 37°C. Plates are then washed with assay buffer and 2,2'-azino-bis(3-ethylbenzothiazoline-6 sulfonic acid) diammonium salt substrate solution is added to each well. The reaction is stopped after 20 min and bound ICAM-1 is determined by measuring the optical density at 405 nm in a microplate reader.

In this assay, compounds of formula I inhibit adhesion of LFA-1 to ICAM-1 with an $\text{IC}_{50} \leq 30 \mu\text{M}$, preferably 0.05 to 30 μM .

ii) Human Mixed Lymphocyte Reaction (MLR)

Peripheral blood mononuclear cells (PBMC) are isolated from human buffy coats. In each experiment, PBMC from three different donors (A, B, and C) are set up in three individual 2-way reactions (A-B, A-C, B-C). Cells are cocultured for six days and proliferation is determined by pulsing the cells with ^3H -thymidine. The concentration of compounds of formula I which results in 50% inhibition of cell proliferation (IC_{50}) is calculated. In this assay, compounds of formula I inhibit the MLR with an IC_{50} in the range of 0.2 to 4 μM .

B. In vivo

i) Murine Thioglycollate Induced Peritonitis

Thioglycollate is injected i.p. to mice and immediately thereafter the compound to be tested is given s.c.. The mice are killed after 4 hours, the peritoneal cavity lavaged and total number of neutrophils in the lavage fluid is determined.

In this assay, the compounds of formula I inhibit thioglycollate induced neutrophil migration when administered s.c. at a dose of from 0.001-50 $\mu\text{g/kg}$.

ii) Allergic Contact Dermatitis (ACD)

Groups of oxazolone-sensitized mice are challenged with 10 μ l of 0.2 or 2.0% oxazolone on the inner surface of the right to elicit ACD. The low concentration of oxazolone is used for testing compounds on systemic activity whereas the high concentration is applied for topical testing. The unchallenged left ears serve as normal controls and dermatitis is evaluated from the individual differences in pinna weight, which is taken as a measure of increase in inflammatory swelling 24 h after the challenge. Dermatitis is evaluated in test- and for comparison in control groups. The test groups are treated with the test compounds either orally (twice, 2 h and immediately before challenge), subcutaneously (immediately before challenge) or topically (30 min after challenge at the site of elicitation of the ACD); the controls are treated similarly with the vehicles alone. For oral and subcutaneous administration the compounds are administered in an oil in water emulsion, for topical administration the compounds are prepared in a mixture of ethanol, acetone and dimethylacetamide. The data of the test- and the vehicle-treated control groups are statistically analysed by ANOVA followed by Dunnet T-test (normal distribution or data) or by H and U-test, respectively. When administered p.o. at a dose of from 0.1 to 10 mg/kg, compounds of formula I inhibit the elicitation phase of allergic contact dermatitis.

iii) Transplantation: Heterotopic mouse heart allograft

The strain combination used: BALB/c \Rightarrow C3H (H-2d \Rightarrow H-2k) comprises MHC and non-MHC mismatch. Female animals are anaesthetised using inhalational isoflurane. Following heparinisation of the donor BALB/c mouse through the abdominal inferior vena cava with simultaneous exsanguination via the aorta, the chest is opened and the heart rapidly cooled. The aorta is ligated and divided distal to the first branch and the brachiocephalic trunk is divided at the first bifurcation. The left pulmonary artery is ligated and divided and the right side divided but left open. All other vessels are dissected free, ligated and divided and the donor heart is removed into iced saline.

The recipient C3H is prepared by dissection and cross-clamping of the infra-renal abdominal aorta and vena cava. The graft is implanted with end-to-side anastomoses, using 11/0 monofilament suture, between the donor brachiocephalic trunk and the recipient aorta and the donor right pulmonary artery to the recipient vena cava. The clamps are removed, the graft tethered retroabdominally, the abdominal contents washed with warm saline and the animal is closed and allowed to recover under a heating lamp. Graft survival is

monitored by daily palpation of the beating donor heart through the abdominal wall. Rejection is considered to be complete when heart beat stops. Improvements of graft function are obtained in animals treated with a compound of formula I administered orally at a daily dose of 30 mg/kg. Significant improvement is obtained when the compound of formula I is administered with an immunosuppressive agent, e.g. cyclosporin A, at a daily dose of 10 mg/kg.

The compounds of formula I are, therefore, useful in the treatment and/or prevention of diseases or disorders mediated by LFA-1/ICAM-1 or ICAM-3 interactions e.g. ischemia/reperfusion injury e.g. myocardial infarction, stroke, gut ischemia, renal failure or hemorrhage shock, acute or chronic rejection of organ or tissue allo- or xenografts, e.g. heart, lung, combined heart-lung, kidney, liver, bowel, bone marrow or pancreatic islets, infection diseases such as septic shock, adult respiratory distress syndrome, or traumatic shock. The compounds of formula I are also useful in the treatment and/or prevention of acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, systemic lupus erythematosus, hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes type I and uveitis, cutaneous manifestations of immunologically-mediated illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, alopecia areata, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitises, seborrhoeic dermatitis, lichen planus, pemphigus, bullous pemphigoid, epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythema multiforme, cutaneous eosinophilias, lupus erythematosus, acne, granuloma annulare, pyoderma gangrenosum, sun burns or toxic epidermal necrolysis), inflammatory bowel disease, ophthalmic inflammatory diseases or immune-mediated conditions of the eye, such as auto-immune diseases, e.g. keratoplasty and chronic keratitis, allergic conditions, e.g. vernal conjunctivitis, inflammatory conditions and corneal transplants. Compounds of formula I are useful as immunosuppressive agents.

For the above uses the required dosage will of course vary depending on the mode of administration, the particular condition to be treated and the effect desired. In general, satisfactory results are indicated to be obtained systemically at daily dosages of from about 0.1 to about 10 mg/kg body weight. An indicated daily dosage in the larger mammal is in the

range from about 0.5 mg to about 80 mg, conveniently administered, for example, in divided doses up to four times a day or in retard form.

For topical use satisfactory results are obtained with local administration of a 1-3 % concentration of active substance several times daily, e.g. 2 to 5 times daily.

The compounds of formula I may be administered systemically or topically, by any conventional route, in particular enterally, e.g. orally, e.g. in the form of tablets or capsules, topically, e.g. in the form of lotions, gels, ointments or creams, or in a nasal or a suppository form. Percutaneous administration via patches or other delivery systems may also be a possible route for prevention or treatment of above diseases.

Pharmaceutical compositions comprising a compound of formula I in association with at least one pharmaceutical acceptable carrier or diluent may be manufactured in conventional manner by mixing with a pharmaceutically acceptable carrier or diluent. Unit dosage forms contain, for example, from about 0.1 mg to about 40 mg of active substance.

Topical administration is e.g. to the skin. A further form of topical administration is to the eye.

The compounds of formula I may be administered in free form or in pharmaceutically acceptable salt form e.g. as indicated above. Such salts may be prepared in conventional manner and exhibit the same order of activity as the free compounds.

In accordance with the foregoing the present invention further provides:

- 1.1 A method for preventing or treating disorders or diseases mediated by LFA-1/ICAM-1 interactions, e.g. such as indicated above, in a subject in need of such treatment, which method comprises administering to said subject an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof;
- 1.2 A method for preventing or treating acute or chronic inflammatory diseases or disorders or autoimmune diseases, e.g. as indicated above, in a subject in need of

such treatment, which method comprises administering to said subject an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof;

2. A compound of formula I, in free form or in a pharmaceutically acceptable salt form for use as a pharmaceutical, e.g. in any of the methods as indicated under 1.1 and 1.2 above.
3. A pharmaceutical composition for use in any of the methods as in 1.1 and 1.2 above comprising a compound of formula I in free form or pharmaceutically acceptable salt form in association with a pharmaceutically acceptable diluent or carrier therefor.
4. A compound of formula I or a pharmaceutically acceptable salt thereof for use in the preparation of a pharmaceutical composition for use in any of the method as in 1.1 and 1.2 above.

The compounds of formula I may be administered as the sole active ingredient or in conjunction with, e.g. as an adjuvant to, other drugs in immunomodulating regimens or other anti-inflammatory agents for the treatment or prevention of allo- or xenograft acute or chronic rejection or inflammatory or autoimmune disorders. For example, the compounds of formula I may be used in combination with cyclosporins, rapamycins or ascomycins, or their immunosuppressive analogs, e.g. cyclosporin A, cyclosporin G, FK-506, ABT-281, ASM981, rapamycin, 40-O-(2-hydroxy)ethyl-rapamycin etc.; corticosteroids; cyclophosphamide; azathioprene; methotrexate; FTY720; leflunomide; mizoribine; mycophenolic acid; mycophenolate mofetil; 15-deoxyspergualine; immunosuppressive monoclonal antibodies, e.g., monoclonal antibodies to leukocyte receptors, e.g., MHC, CD2, CD3, CD4, CD7, CD25, CD28, B7, CD40, CD45 or CD58 or their ligands; or other immunomodulatory compounds, e.g. CTLA4Ig, or other adhesion molecule inhibitors, e.g. mAbs or low molecular weight inhibitors including Selectin antagonists and VLA-4 antagonists. A preferred composition is with Cyclosporin A, FK506, rapamycin or 40-O-(2-hydroxy)ethyl-rapamycin.

Where the compounds of formula I are administered in conjunction with other immunosuppressive / immunomodulatory or anti-inflammatory therapy, e.g. for preventing or

treating chronic rejection as hereinabove specified, dosages of the co-administered immunosuppressant, immunomodulatory or anti-inflammatory compound will of course vary depending on the type of co-drug employed, e.g. whether it is a steroid or a cyclosporin, on the specific drug employed, on the condition being treated and so forth. In accordance with the foregoing the present invention provides in a yet further aspect:


5. A method as defined above comprising co-administration, e.g. concomitantly or in sequence, of a therapeutically effective amount of a compound of formula I in free form or in pharmaceutically acceptable salt form, and a second drug substance, said second drug substance being an immunosuppressant, immunomodulatory or anti-inflammatory drug, e.g. as indicated above.
6. A therapeutic combination, e.g. a kit, for use in any method as defined under 1.1 or 1.2 above, comprising a compound of formula I, in free form or in pharmaceutically acceptable salt form, with at least one pharmaceutical composition comprising an immunosuppressant, immunomodulatory or anti-inflammatory drug. The kit may comprise instructions for its administration.

Compounds of Examples 2, 28 and 34 are preferred, particularly for use in the treatment of inflammatory skin diseases, e.g. as indicated above. In one test run, following results were obtained: an IC_{50} of 0.05, 0.79 and 0.19 μM , respectively for the compounds of Ex. 2, 28 and 34, in the test Ai); an IC_{50} of 0.2 μM for the compound of Ex. 2 in the MLR test Aii); an ED_{50} of 0.1 $\mu g/kg$ p.o. for the compound of Ex. 2 in the test Bi); in Bii) compound of Ex. 2 has an inhibiting effect of 41% when administered p.o. at a dose of 2 x 3 mg/kg and compound of Ex. 34 inhibits inflammatory swelling by 41% at 2x 1 mg/kg p.o.

Preferred compounds of formula I are those inhibiting HMG CoA Reductase activity with an $IC_{50} \geq 1 \mu M$, e.g. $\geq 50 \mu M$, in the in vitro Microsomal assay as disclosed in WO 99/11258.

CLAIMS

1. A compound of formula I



wherein

each of a—b and α—β independently, is either a single bond or a double bond;


R₁ is



wherein R_a is H, C₁₋₆alkyl optionally substituted by OH or C₁₋₄alkoxy, C₂₋₆alkenyl or aryl-C₁₋₄alkyl;

R₂ is OH; -O-CO-R₅ wherein R₅ is C₁₋₈alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkyl, aryl or aryl-C₁₋₄alkyl; or -O-R₆ wherein R₆ is the residue of an α-amino-acid attached to O through its carbonyl residue or -CHR₇-COR₈ wherein R₇ is H, C₁₋₄alkyl, heteroC₁₋₄alkyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkyl, aryl or aryl-C₁₋₄alkyl and R₈ is OH, C₁₋₄alkoxy or NR₉R₁₀ wherein each of R₉ and R₁₀ independently is H, C₁₋₄alkyl or R₉ and R₁₀ form together with the nitrogen to which they are bound, a heteroaryl group;

R₃ is a substituted lactam, piperidyl, linear amino alcohol or cyclic carbamate, or a residue of formula (i)



wherein

R₁₃ is OH; C₁₋₆alkoxy; -O-CO-C₁₋₆alkyl; or -O-CO-NHC₁₋₆alkyl;

R₁₄ is OH; C₁₋₄alkoxy; C₁₋₄alkyl; C₁₋₄alkoxy-carbonyl-C₁₋₄alkoxy; hydroxy-C₁₋₅alkoxy;

C₁₋₄alkoxy-C₁₋₅alkoxy; C₁₋₄alkoxy-carbonyl-C₁₋₄alkyl; or NR_{9a}R_{10a}-C₁₋₅alkoxy wherein each of R_{9a} and R_{10a} independently has one of the significances given for R₉ and R₁₀;




R_{15} is H or C_{1-4} alkyl; and

R_{16} is $CONR_{17}R_{18}$ wherein one of R_{17} and R_{18} is H and the other is C_{1-6} alkyl, hydroxy- C_{1-6} alkyl, C_{3-7} cycloalkyl- C_{1-4} alkyl or aryl- C_{1-4} alkyl; or C_{1-6} alkoxy-carbonyl; each of α — β and α — β being a single bond when each of R_{13} or R_{14} is OH; and


R_4 is H or OR_{19} wherein R_{19} is C_{1-6} alkyl, hydroxy- C_{1-6} alkyl, C_{1-4} alkoxy- C_{1-6} alkyl, aryl- C_{1-4} alkyl or C_{1-4} alkoxycarbonyl- C_{1-4} alkyl,

and wherever "aryl" appears as is or in the significances of "aryl- C_{1-4} alkyl" in the above definition, it is "phenyl" or "naphthyl" optionally substituted by halogen, OH, $NR_{11}R_{12}$, COOH, CF_3 , C_{1-4} alkoxy, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, hydroxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl, cyano or $CONR_{11}R_{12}$, each of R_{11} and R_{12} independently being H, C_{1-4} alkyl, phenyl, naphthyl, phenyl- C_{1-4} alkyl or naphthyl- C_{1-4} alkyl or R_{11} and R_{12} together with the nitrogen to which they are bound forming heteroaryl; and wherever "heteroaryl" appears, it is a 5- or 6-membered heterocyclic residue optionally fused to a benzene ring; in free form or in salt form.

2. A compound according to claim 1 wherein R_3 is a radical of formula (a), (b), (c₁) or (c₂)



or



wherein

R_{30} is C_{1-8} alkyl; C_{3-7} cycloalkyl; aryl; C_{3-7} cycloalkyl- C_{1-4} alkyl; aryl- C_{1-4} alkyl; heteroaryl; or heteroaryl- C_{1-4} alkyl;

R_{31} is OH; C_{1-4} alkoxy; C_{1-4} alkyl; C_{1-4} alkoxy-carbonyl- C_{1-4} alkoxy; hydroxy- C_{1-5} alkoxy; C_{1-4} alkoxy- C_{1-5} alkoxy; C_{1-4} alkoxy-carbonyl- C_{1-4} alkyl; amino- C_{1-4} alkoxy; $HOOC-C_{1-4}$ alkoxy; $HOOC-C_{1-4}$ alkyl; or $NR_{9a}R_{10a}-C_{1-5}$ alkoxy wherein each of R_{9a} and R_{10a} independently is H, C_{1-4} alkyl or R_{9a} and R_{10a} form together with the nitrogen to which they are bound, a heteroaryl group;

R_{40} has one of the significances given for R_{30} ;

R_{41} has one of the significances given for R_{31} or is $-O-CO-C_{1-8}$ alkyl;

either each of X and Y is H or X and Y form together



each of R_{50} , independently is H; C_{1-8} alkyl; C_{3-7} cycloalkyl; aryl; C_{3-7} cycloalkyl- C_{1-4} alkyl; aryl- C_{1-4} alkyl; heteroaryl; heteroaryl- C_{1-4} alkyl; C_{1-4} alkyl-carbonyl; aryl-carbonyl; heteroaryl-carbonyl; aryl- C_{1-4} alkyl-carbonyl or heteroaryl- C_{1-4} alkyl-carbonyl, and

each of R_{51} , independently is H; C_{1-4} alkyl; hydroxy- C_{1-4} alkyl; amino- C_{1-4} alkyl; C_{1-4} alkoxy- C_{1-4} alkyl; C_{1-4} alkoxy-carbonyl- C_{1-4} alkyl wherein C_{1-4} alkoxy is optionally substituted by amino, C_{1-4} alkyl-amino or di- $(C_{1-4}$ alkyl)-amino; $HOOC-C_{1-4}$ alkyl; or $R_{23}R_{24}N-CO-C_{1-4}$ alkyl wherein R_{23} is H, C_{1-4} alkyl, hydroxy- C_{1-4} alkyl, polyhydroxy- C_{1-8} alkyl, heteroaryl, heteroaryl- C_{1-4} alkyl, amino- C_{1-4} alkyl, C_{1-4} alkylamino- C_{1-4} alkyl, di- $(C_{1-4}$ alkyl)amino- C_{1-4} alkyl or aryl- C_{1-4} alkyl and R_{24} is H, C_{1-4} alkyl or hydroxy- C_{1-4} alkyl,

at least one of R_{50} and R_{51} being other than H.

3. A compound according to claim 2 wherein R_3 is a radical of formula (a);

R_{30} being benzyl or naphthyl-methyl wherein the phenyl or naphthyl ring is optionally substituted by OH, C_{1-4} alkoxy, hydroxy- C_{1-4} alkoxy or hydroxy- C_{1-4} alkyl, or morpholino, pyridyl, indolyl or quinolyl; and

R_{31} being OH, C_{1-4} alkoxy, hydroxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl- C_{1-4} alkoxy or $HOOC-C_{1-4}$ alkoxy.

4. A compound according to claim 2 wherein R_3 is a radical of formula (c₁) or (c₂) wherein X and Y form together $-CO-$;

R_{50} being benzyl or naphthyl-methyl wherein the phenyl or naphthyl ring is optionally substituted by OH, C_{1-4} alkoxy, hydroxy- C_{1-4} alkoxy or hydroxy- C_{1-4} alkyl; and

- R_{51} being hydroxy- C_{1-4} alkyl; amino- C_{1-4} alkyl; C_{1-4} alkoxy- C_{1-4} alkyl; C_{1-4} alkoxy-carbonyl- C_{1-4} alkyl; $HOOC-C_{1-4}$ alkyl; or $R_{23}R_{24}N-CO-C_{1-4}$ alkyl wherein R_{23} and R_{24} are as defined in claim 2.
5. A compound according to claim 1 which is (S)-2-methyl-butyric acid (S)-(3R,7S,8aR)-8-((S)-2-[(4R,6R)-3-(4-hydroxy-3-methoxy-benzyl)-4-methylcarbamoylmethyl-2-oxo-[1,3]oxazinan-6-yl]-ethyl)-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester or (S)-2-methyl-butyric acid (1S,3R,7S,8S,8aR)-8-{2-[(2S,4R)-4-hydroxy-1-(5-hydroxymethyl-6-methoxy-naphthalen-2-yl)methyl]-6-oxo-piperidin-2-yl]-ethyl}-3,7-dimethyl-1,2,3,7,8,8a-hexahydro-naphthalen-1-yl ester.
 6. A compound according to any one of claims 1 to 5, in free form or in pharmaceutically acceptable salt form, for use as a pharmaceutical.
 7. A pharmaceutical composition comprising a compound of formula I according to any one of claims 1 to 5, in free form or in pharmaceutically acceptable salt form, in association with a pharmaceutically acceptable diluent or carrier therefor.
 8. A pharmaceutical composition according to claim 7, for use in combination with at least one pharmaceutical composition comprising an immunosuppressant, immunomodulatory or anti-inflammatory drug.
 9. A method for preventing or treating disorders or diseases mediated by LFA-1/ICAM-1 interactions in a subject in need of such treatment, which method comprises administering to said subject an effective amount of a compound of formula I according to any one of claims 1 to 5 or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/01191

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C235/26 C07C237/18 C07C235/30 C07C237/20 C07D211/76
C07D265/24 C07D401/06 A61K31/4402 A61K31/45 A61K31/4545
A61K31/535 A61P9/10 A61P3/06 A61P29/00 A61P37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 99 11258 A (NOVARTIS AG ET AL.) 11 March 1999 (1999-03-11) cited in the application the whole document	1,7
Y	WO 97 16184 A (WARNER-LAMBERT COMPANY) 9 May 1997 (1997-05-09) the whole document	1,7
Y	EP 0 415 488 A (MERCK & CO. INC) 6 March 1991 (1991-03-06) the whole document	1,7
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

26 June 2000

Date of mailing of the international search report

30/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 851 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01191

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61P43/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KLAUS WENKE ET AL.: "Simvastatin reduces graft vessel disease and mortality after heart transplantation" CIRCULATION, vol. 96, no. 5, 2 September 1997 (1997-09-02), pages 1398-1402, XP002090500 the whole document	1,7
Y	SATORU NIWA ET AL.: "Inhibitory effect of fluvastatin, an HMG-CoA Reductase inhibitor. On the expression of adhesion molecules on human monocyte cell line." INT. J. IMMUNOPHARMAC., vol. 18, no. 11, 1996, pages 669-675, XP002090501 the whole document	1,7

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

26 June 2000

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kyriakakou, G

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/01191

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>P. DINAPOLI ET AL.: "Does 3-Hydroxy-3-methylglutaryl Coenzyme A Reductase Inhibitor Therapy Exert a Direct Anti-Ischemic Effect?" CIRCULATION, vol. 97, no. 9, 10 March 1998 (1998-03-10), page 937 XP002090499 dallas, tx the whole document</p>	1,7
Y	<p>BRUNO REICHART ET AL.: "What is the role of lipid lowering therapy in heart-allograft failure?" KIDNEY INTERNATIONAL, vol. 48, no. suppl.52, December 1995 (1995-12), pages s52-s55, XP002090502 the whole document</p>	1,7

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 00/01191

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9911258 A	11-03-1999	AU 9739198 A	22-03-1999
		EP 1007033 A	14-06-2000
		ZA 9807787 A	01-03-1999
WO 9716184 A	09-05-1997	AU 7253996 A	22-05-1997
		BG 102417 A	29-01-1999
		BR 9611410 A	05-01-1999
		CA 2233558 A	09-05-1997
		CN 1201389 A	09-12-1998
		CZ 9801271 A	16-12-1998
		EP 0858336 A	19-08-1998
		HU 9901865 A	28-10-1999
		JP 11515025 T	21-12-1999
		NO 981961 A	04-05-1998
		PL 326365 A	14-09-1998
		SK 55798 A	11-06-1999
EP 415488 A	06-03-1991	CA 2024248 A	01-03-1991
		DE 69008277 D	26-05-1994
		JP 3184940 A	12-08-1991
		US 5098931 A	24-03-1992